

## When do you step up or increase long-term control medications?

To regain control when the child has:

- Increased symptoms and/or nighttime awakenings due to symptoms.
- Reduced ability to play, exercise, or participate in normal daily activities.
- Daily or increasing use of short-acting  $\beta_2$ -agonists.
- Reduction in PEF (by about 20%).
- When the goals of therapy are not being achieved.

## How to help the child regain control of asthma:

- A short (3- to 10-day) course of oral corticosteroids may be needed to speed the resolution of moderate or severe exacerbations, or to reestablish control during a period of gradually deteriorating symptoms.
- If a short course of oral corticosteroids does not control symptoms, is effective for only a short time (i.e., < 2 weeks), or if the child used oral corticosteroids frequently, consider a step-up or increase in the long-term control medication. This can be achieved by:
  - ⇒ Increasing the dose.
  - ⇒ Increasing the frequency of dosing.
  - ⇒ Using or adding a different medication.



## Before increasing medications, assess possible reasons for poor asthma control:

- Inhaler technique
- Adherence
- Environmental exposures
- Complicating factors (e.g., upper respiratory infections)

## Remember the goals of therapy:

- Prevent chronic and troublesome symptoms.
- Prevent exacerbations of symptoms.
  - ⇒ No acute episodes of asthma that require a doctor visit, emergency room visit, or hospital stay.
- Maintain normal activity levels.
- Maintain “normal” pulmonary function.
- Minimal (ideally no) adverse effects from medication.

## **Children with severe persistent asthma may need to use oral corticosteroids on a long-term basis.**

Children with asthma whose symptoms are not controlled on high doses of inhaled corticosteroids *and* long-acting bronchodilators may need to use oral corticosteroids on a regular, long-term basis. For these children:

- Use the lowest possible dose (single dose, daily or on alternate days).
- Monitor the child closely for corticosteroid adverse effects.
- Make persistent attempts to reduce use of oral corticosteroids when control of asthma is achieved. High doses of inhaled corticosteroids are preferable because of fewer adverse effects.
- Regular consultation with an asthma specialist is recommended.

### **Remember when using the stepwise approach to therapy that referral to an asthma specialist for consultation or co-management is recommended when:**

- The child has had a life-threatening asthma exacerbation.
- Goals of asthma therapy are not being met after 3-6 months of treatment; earlier if the child appears unresponsive to treatment.
- Signs and symptoms are atypical, or there are problems in differential diagnosis.
- Other conditions complicate asthma or its diagnosis (e.g., untreated sinusitis, rhinitis).
- Additional diagnostic testing is indicated (e.g., pulmonary function testing, allergy skin testing).
- The child or family needs additional education and guidance on complications of therapy, problems with adherence, or avoidance of triggers.
- The child is being considered for immunotherapy.
- The child has severe persistent asthma.
- The child is under 3 years of age and has moderate or severe persistent asthma.
- The child has used long-term oral corticosteroid therapy, high-dose inhaled corticosteroid therapy, or more than 2 bursts of oral corticosteroids in 12 months.

## **Treat asthma exacerbations promptly and aggressively.**

- All children with asthma need an inhaled, short-acting  $\beta_2$ -agonist (MDI, nebulizer, DPI) for exacerbations.
- Give the child, the child's family, and caregivers (daycare providers, teachers, coaches, scout leaders, camp counselors, school and camp nurses) a written, easy-to-understand ACTION PLAN to manage exacerbations. Include:
  - ⇒ The early signs of worsening asthma.
  - ⇒ Which medications to use and how to use them.
  - ⇒ Specific instructions for when to contact the physician or emergency room.
- A short (3- to 10-day) course of oral corticosteroids may be needed to speed the resolution of moderate persistent or severe persistent exacerbations, or to reestablish control during a period of gradually deteriorating symptoms.
- Exacerbations of asthma symptoms (coughing, wheezing, shortness of breath or rapid breathing, chest tightness) with viral respiratory infections are common in children, and need to be treated appropriately with adequate doses of short-acting  $\beta_2$ -agonists and, in some cases, oral corticosteroids.
- Asthma exacerbations may require treatment at the physician's office or emergency room.
  - ⇒ Frequent short-acting  $\beta_2$ -agonist, by nebulization.
  - ⇒ Consider adding ipratropium bromide.
  - ⇒ Consider oxygen to relieve hypoxemia.
  - ⇒ Pulse oximetry is recommended to follow oxygen saturation.
  - ⇒ Monitor the response to therapy with serial measurements of FEV<sub>1</sub> or PEF.

**All children with asthma should have a short-acting bronchodilator to provide prompt relief of acute symptoms (coughing, wheezing, difficulty breathing, chest tightness).**

**Over-the-counter MDIs are not as effective as short-acting, inhaled  $\beta_2$ -agonists, and may unnecessarily delay seeking medical care.**

**All offices that treat acute exacerbations of asthma should have a peak flow meter, a nebulizer, an oximeter, and oxygen.**

# Assessing the Severity of an Asthma Exacerbation

**Asthma severity is a continuum. Any child, regardless of overall severity, can have a severe exacerbation.**

## Risk Factors for Death from Asthma

- Past history of sudden severe exacerbations
- Prior intubation for asthma
- Prior admission to intensive care unit for asthma
- $\geq 2$  hospitalizations for asthma in past 12 months
- $\geq 3$  emergency care visits for asthma in past 12 months
- Hospitalization or emergency care visit for asthma in past month
- Use of  $> 1$  canister/month of inhaled short-acting  $\beta_2$ -agonist
- Current chronic use of oral corticosteroids
- Difficulty perceiving airflow obstruction or its severity
- Low socioeconomic status and urban residence
- Illicit drug use
- Serious psychiatric disease or psychosocial problems

## Look at the child for signs of distress:

- **Shortness of breath or rapid breathing**
  - ⇒ As severity increases, the child may have difficulty talking or laying down. The infant may have difficulty feeding and may have a shorter, softer cry.
- **Increased respiratory rate** (see table below)
- **Use of accessory muscles with retractions**
- **Wheezing**
  - ⇒ In mild exacerbations, wheezing is evident on expiration.
  - ⇒ As the severity of the exacerbation increases, both inspiratory and expiratory wheezing may be present.
  - ⇒ During a severe exacerbation, the chest may be “silent.”
- **Decreased PEF**
- **Agitation**

## Normal Breathing and Pulse Rates for Children


Age	Breathing rate (awake)	Pulse rate
< 2 months	< 60/minute	< 160/minute
2-12 months	< 50/minute	< 120/minute
12-24 months	< 40/minute	< 110/minute
2-5 years	< 40/minute	< 110/minute
6-8 years	< 30/minute	< 110/minute

# Managing Asthma Exacerbations in the Home

## Assess severity

**Measure PEF:** < 50% predicted or personal best suggests severe exacerbation.

**Note signs and symptoms:** Degree of cough, shortness of breath, wheeze, chest tightness correlate imperfectly with severity. Accessory muscle use and retractions (sucking in of chest) suggest severe exacerbation.



## Initial treatment


Inhaled short-acting beta<sub>2</sub>-agonist: up to 3 Tx of 2-4 puffs by MDI every 20 min, or 1 nebulizer Tx



Response to treatment	Good	Incomplete	Poor
<b>Exacerbation</b> PEF (predicted or personal best) Wheezing or shortness of breath	<i>Mild</i> > 80% None	<i>Moderate</i> 50%-80% Persistent	<i>Severe</i> < 50% Marked
<b>Continued Treatment</b>	Continue inhaled, short-acting beta <sub>2</sub> -agonist every 3-4 hrs for 24-48 hrs. For children on inhaled corticosteroids, double dose for 7-10 days.	Continue inhaled short-acting beta <sub>2</sub> -agonist. Add oral corticosteroid.	Repeat beta <sub>2</sub> -agonist immediately. Add oral corticosteroid.
<b>Follow-up</b>	Contact clinician.	Contact clinician same day.	If distress is severe and child is non-responsive, CALL DOCTOR IMMEDIATELY, AND PROCEED TO EMERGENCY DEPARTMENT. CONSIDER CALLING AMBULANCE OR 911.



# Management of Asthma Exacerbations: Emergency Department and Hospital-Based Care



# Special Considerations when Treating Children with Asthma

## Administering asthma medications to infants and children can be challenging.

- Medications to treat asthma in children may be given by inhalation or orally (as tablets or liquids).
  - ⇒ The inhaled route is generally preferred because:
    - Higher concentrations can be delivered more effectively to the airways.
    - Systemic side effects are avoided or minimized.
    - The onset of action of short-acting  $\beta_2$ -agonists is substantially shorter when inhaled.
- Dosages reaching the airway will vary considerably depending on the route of administration (and the device(s) used for inhaled medications).
- The ability of individual children to use different devices for inhaled medications may vary considerably.

## For children < 2 years:

- Nebulizer therapy with face mask may be preferred for administering cromolyn sodium, and for short-acting  $\beta_2$ -agonists during exacerbations.
- Short-acting  $\beta_2$ -agonists are available as liquids, but:
  - ⇒ The onset of action is slower than when given by inhalation (approximately 30 minutes for the liquid as compared to several minutes for inhalation).
  - ⇒ Adverse effects (e.g., tremor, irritability) are more likely.
- Drugs administered by MDI may be given using a spacer/holding chamber and face mask.
  - ⇒ The dose will be variable.
  - ⇒ Inhaled corticosteroids by MDI should always be given with a spacer/holding chamber.

## How do you give medication to infants and young children?

- Use a metered dose inhaler with a spacer/holding chamber plus face mask.
- Use a nebulizer.
- Use a liquid.

*Liquid albuterol does not have as quick an onset-of-action as inhaled albuterol, a consideration for treating exacerbations.*

**Tailor the delivery device to the child, taking the child's needs into consideration.**

### **For children between 3 and 5 years:**

- Inhaled medications are preferred.
- Some children can use an MDI and spacer/holding chamber.
- If the desired therapeutic effects are not achieved, or if the child cannot use an MDI with spacer/holding chamber, a nebulizer or MDI plus spacer/holding chamber with face mask may be required.

### **For school-age children:**

- Inhaled medications are preferred.
- MDIs, DPIs, and nebulizers may be used.
- The child should be able to produce the necessary effort and coordination needed for the specific device.
- All inhaled corticosteroids by MDI should be used with a spacer/holding chamber.
- Some children carry their short-acting beta<sub>2</sub>-agonist MDI without a spacer/holding chamber. This is acceptable if the child has demonstrated good technique.
- School medication policies should be known.





## Types of Inhalation Devices for Asthma Medications

Device/Medications	Age <sup>1</sup>	Comments
<b>Metered-dose inhaler (MDI)</b> Beta <sub>2</sub> -agonists Corticosteroids Cromolyn sodium Nedocromil sodium Anticholinergics	> 5 years (< 5 years with spacer/holding chamber and face mask for some children)	The child may have difficulty triggering a puff while inhaling. Use with a spacer/holding chamber helps.
<b>Breath-actuated MDI</b> Beta <sub>2</sub> -agonists	> 5 years	The child may not be able to generate the necessary inspiratory flow. Device does not require the use of holding chamber or spacer.
<b>Dry-powder inhaler (DPI)</b> Beta <sub>2</sub> -agonists Corticosteroids	> 5 years (can be used in 4 year olds, but delivery is more consistent over 5)	Some devices deliver drug more effectively than an MDI. Some devices may not work in children with low inspiratory volumes.
<b>Nebulizer</b> Beta <sub>2</sub> -agonists Cromolyn sodium Anticholinergics	Patients of any age who cannot use an MDI with spacer/holding chamber or with face mask.	Useful in infants and very young children, and any child with a moderate to severe asthma episode. Delivery method of choice for cromolyn sodium.

<sup>1</sup>These ages are suggested as guides for making clinical decisions. The clinician must use his/her judgment to tailor treatment to the specific needs and circumstances of the child and family.

### Using an MDI with a spacer/holding chamber may be easier than using an MDI alone.

- Trigger 1 puff from MDI into spacer/holding chamber for each inhalation.
- Some young children may be able to use an MDI with spacer/holding chamber and face mask.
  - ⇒ If a face mask is used, allow 3 to 5 inhalations for each puff triggered from the MDI.

### Advantages of using spacers and holding chambers:

- Many children have a difficult time inhaling while pressing the inhaler to trigger a puff. Simple tubes do not take care of this problem!
- Decreases oropharyngeal deposition.
- Reduces possible side effects, particularly with inhaled corticosteroids.
  - ⇒ Potential side effects of inhaled corticosteroids include oral candida (thrush), dysphonia, and reflex cough and bronchospasm (also evident with other inhaled medications).

## The child's schedule and giving asthma medications

**Instruct the child and parent in the appropriate use of all medications and devices. Advise the parent to share this information with ALL caregivers.**

**Have the child (and parent) bring and demonstrate use of their delivery devices (including inhalers, spacer/ holding chambers, face masks, nebulizers) at each office visit.**

- Provide an action plan for handling exacerbations, including the clinician's recommendation regarding self-administration of medication and plans to ensure prompt, reliable access to medications.
- If possible, schedule long-term control medications so that they are not taken at school, even if this results in uneven dosing intervals.
  - ⇒ However, some children benefit from close supervision of therapy. For these children, giving medication at school, under the supervision of a school health professional, is recommended.
- It may be helpful for some younger children to have a compressor-driven nebulizer available at their school or daycare facility.

### **Reliable, prompt access to asthma medication is essential during the day.**

- The child, caregivers, teachers, school nurses, and school boards should understand this.
- Older children should be allowed to carry and self-administer quick-relief medications, with physician and parent approval.



## Infants and adolescents have special needs.

### 0 to 18 months

- **Early management of wheezing** may alter the course of asthma later on. However, infants can be difficult to manage. Referral to a pediatric asthma specialist for consultation or co-management should be considered for children being given daily medication.
- **Carefully monitor the response to medication.**
  - ⇒ Treat the wheeze for 4 to 8 weeks.
  - ⇒ If there is no clear response, stop treatment.
  - ⇒ Consider alternate therapy or an alternate diagnosis if the child is not: growing and developing normally, eating normally and gaining weight, or sleeping.
- **Diet is usually not a factor** for the wheezing child. (When there are reactions after ingesting food, the most common causes are milk, peanut products, soy, wheat, eggs.)
- **Reducing exposure to viral respiratory infections may be important** for the difficult-to-manage wheezer.



### Many children have multiple caretakers.

All homes and caretakers should have medications, devices, and a management plan for how and when to use them.

### Upper respiratory viral infections are a key precipitating factor of asthma symptoms in young children.

### Primary prevention of asthma (preventing initial development) may alter its course.

- Minimize exposure to dust mites, tobacco smoke, animal dander, cockroach allergens.
  - ⇒ Exposure to high levels of dust mite and tobacco smoke are associated with an increased incidence of asthma among infants.
- Prolonged breast feeding and avoiding the early introduction of allergenic foods may reduce eczema and food sensitization, but have not been shown to reduce the prevalence of asthma.

### **Special concerns of the adolescent include:**

- Effects of medications on appearance (height, weight, acne), ability to exercise, menses.
- Taking medications in public.
- Peer pressure.

### **Poorly controlled asthma may delay growth.**

### **The potential risks of inhaled corticosteroids are well balanced by their benefits.**

## **Adolescents**

- LISTEN to the patient! Find out their expectations and goals.
- Find out what the adolescent is willing to do, and then work out a management plan together.
- Ask the adolescent about smoking, and exposure to tobacco smoke and possible drug use.
- Encourage exercise and physical activity. Asthma should not be an excuse for not participating in physical education or sports.
  - ⇒ Develop an asthma management plan that will allow them to participate in any activity that they wish.
  - ⇒ Make it easy to take medications before exercise.
- Consider symptoms related to hobbies and workplace exposures.
- Consider nonadherence if the teen is not doing well.

## **Asthma, inhaled corticosteroids, and linear growth**

- Poorly controlled asthma may delay growth.
- Growth rates are highly variable. Short-term evaluations may not predict final adult height.
- Children with asthma tend to have longer periods of reduced growth rates before puberty (males more than females).
- Some studies of the use of inhaled corticosteroids to treat asthma in children have identified growth delay; others have not. The clinical significance of the findings is unclear. Monitoring growth is recommended.
  - ⇒ For children with mild or moderate persistent asthma, medium-dose inhaled corticosteroid therapy may be associated with a possible, but not predictable, adverse effect on linear growth.
  - ⇒ For children with severe persistent asthma, use of high-doses of inhaled corticosteroids has significantly less potential for adverse growth effects than oral corticosteroids.
  - ⇒ Use of spacers will minimize local and systemic effects of inhaled corticosteroids that are delivered by MDI's.

## Preventing “Anticipated” Episodes of Asthma Symptoms

When you understand and recognize the triggers of asthma symptoms for a child, it is possible to prevent or at least minimize episodes of symptoms due to “anticipated exposures” to the trigger.

For “anticipated” symptoms upon exposure to:	Treat with:
Exercise	<ul style="list-style-type: none"><li>• Short-acting, inhaled beta<sub>2</sub>-agonist, cromolyn sodium, or nedocromil sodium shortly (5-30 minutes) before exercising.</li><li>• Long-acting beta<sub>2</sub>-agonist taken at least 30 minutes before exercising.</li><li>• Regular use of a long-term controller medication may reduce the likelihood of exercise-induced symptoms.</li></ul>
Cold (dry) air	<ul style="list-style-type: none"><li>• Short-acting, inhaled beta<sub>2</sub>-agonist, cromolyn sodium, or nedocromil sodium shortly (5-30 minutes) before going out in the cold.</li><li>• Cover nose and mouth with a scarf on cold or windy days.</li></ul>
Allergens (e.g., animal dander, pollens)	<ul style="list-style-type: none"><li>• Cromolyn sodium or nedocromil sodium shortly (5-30 minutes) before anticipated exposure.</li></ul>

## Managing the child with seasonal symptoms

Some children experience asthma symptoms only in relationship to certain pollens and molds. If the child has seasonal asthma on a predictable basis, long-term control medication should be:

- Started (or increased) prior to the anticipated onset of symptoms and continued through the season, and
- Gradually decreased (or stopped) after the season.



**Viral upper respiratory infections are a key precipitating factor of asthma symptoms in young children.**

## **Allergen immunotherapy may be considered for children with asthma when:**

- There is clear evidence of a relationship between symptoms and exposure to an unavoidable allergen to which the child is sensitive.
- Symptoms occur all year or during a major portion of the year.
- There is difficulty controlling symptoms with pharmacologic management because multiple medications are required, medications are ineffective, or medications are not accepted by the child (or parents).

Referral to a specialist for consultation or co-management is recommended for children being considered for allergy immunotherapy. Allergen immunotherapy should be administered only in an office where facilities and trained personnel are available to treat any reaction that may occur. Life threatening reactions can occur, but are rare.

## **Treating asthma symptoms due to viral upper respiratory infections (URIs)**

- **For mild symptoms**, a short-acting, inhaled beta<sub>2</sub>-agonist (every 4 to 6 hours for 24 hours, longer with physician consult) may be sufficient to control the symptoms and to improve lung function.
- **For children with recurrent URIs**, when this therapy needs to be repeated more frequently than every 6 weeks, a seasonal increase in (or starting) long-term control therapy may be necessary.
- **For a moderate persistent or severe persistent exacerbation**, a short (3- to 10-day) course of oral corticosteroids should be considered.
- **If the child has a history of severe recurrent exacerbations with URIs**, consider initiating a short (3- to 10-day) course of oral corticosteroids at the first sign of URI.
- **Appropriate long-term control therapy during the viral season** may reduce the frequency and severity of viral-induced symptoms.

## Managing exercise-induced asthma

- **Exercise-induced bronchospasm (EIB) is caused** by a loss of heat, water, or both from the airways during exercise due to hyperventilation of cool, dry air relative to the air within the lungs.
- **EIB usually begins during exercise and peaks** 5 to 10 minutes after stopping exercise.
- **Exercise may be the only trigger of asthma** for some children, particularly children with allergic rhinitis.
  - ⇒ These children should be monitored regularly because EIB is often a marker of inadequate asthma management.
- **Symptoms may spontaneously resolve** within 1 hour after exercise.
- **A warm-up period** before exercise may help.
- If asthma symptoms occur regularly with usual activities or exercise, increasing (or adding) daily long-term control medications may be warranted.
- **Teachers and coaches** need to be notified that a child has exercise-induced symptoms.
  - ⇒ They should be told that the child can participate in activities, but may need inhaled medication before the activity.
- **Appropriate long-term control therapy**, especially with anti-inflammatory medications, can reduce the frequency and severity of exercise-induced symptoms.

**Exercise-induced symptoms should be anticipated in ALL children with asthma.**

**Asthma should not be an excuse from participating in physical education, sports, or exercise.**

- Develop an asthma management plan that will allow the child to participate in any activity that they wish.
- Make it easy to take medications before exercise.
- If full activity is not possible, have the child participate to the extent that he/she can.

**Sometimes physical education/activities DO need to be adjusted for the child with asthma. For example:**

- On a severe allergy day
- On a windy day
- In extreme cold
- During and/or after a viral URI

Treatment for exercise-induced asthma:	Minutes before exercise to take medication:	May prevent symptoms for up to:
Short-acting inhaled beta <sub>2</sub> -agonist (2-4 puffs)	5-30 (best taken just before exercising)	2-3 hours
Cromolyn sodium or nedocromil sodium (2-4 puffs)	5-30 (best taken just before exercising)	1-2 hours
Long-acting beta <sub>2</sub> -agonist (2 puffs)	At least 30	10-12 hours

## References

- Agertoft L, Pedersen S. Importance of the inhalation device on the effect of budesonide. *Arch Dis Child* 1993; 69:130-133.
- Ahrens R, Lux C, Bahl T, Han SH. Choosing the metered-dose inhaler spacer or holding chamber that matches the patient's need: evidence that the specific drug being delivered is an important consideration. *J Allergy Clin Immunol* 1995; 96:288-294.
- Albazzaz MK, Neale MG, Patel KR. Dose-response study of nebulized nedocromil sodium in exercise induced asthma. *Thorax* 1989; 44:816-819.
- Allen DB, Lemanske RF Jr. The safety of chronic asthma treatments: continuous beta agonist therapy and prolonged inhaled corticosteroids in childhood asthma. In: Middleton E Jr, Reed CE, Ellis EF, Adkinson NF Jr, Yunginger JW, Busse WW, eds. *Allergy: Principles and Practice*. 4<sup>th</sup> ed. St. Louis, MO.: Mosby Yearbook, 1993; 1-16.
- Allen DB, Mullen M, Mullen B. A meta-analysis of the effect of oral and inhaled corticosteroids on growth. *J Allergy Clin Immunol* 1994; 93:967-976.
- Allen DB. Growth suppression by glucocorticoid therapy. In: Vassallo J, ed. *Endocrinology and metabolism clinics in North America*. Philadelphia: W.B. Saunders Co. 1996; 699-717.
- Amirav I, Newhouse MT. Metered-dose inhaler accessory devices in acute asthma: efficacy and comparison with nebulizers: a literature review. *Arch Ped Adolesc Med* 1997; 151:876-882.
- Anderson SD. Issues in exercise-induced asthma. *J Allergy Clin Immunol* 1985; 76:763-772.
- Balfour-Lynn L. Growth and childhood asthma. *Arch Dis Child* 1986; 61(11):1049-1055.
- Barnes PJ, Pedersen S. Efficacy and safety of inhaled corticosteroids in asthma. *Am Rev Respir Dis* 1993; 148:S1-S26.
- Barnes PJ. Current therapies for asthma. Promise and limitations. *Chest* 1997; 111:S17-S26.
- Barnes PJ. From pathophysiological mechanisms to pharmacological treatment of childhood asthma. *Ped Pulmonol* 1995; 11:40-41.
- Barnes PJ. Inhaled glucocorticoids for asthma. *N Engl J Med* 1995; 332:868-875.
- Barros MJ, Rees PJ. Bronchodilator responses to salbutamol followed by ipratropium bromide in partially reversible airflow obstruction. *Respir Med* 1990; 84:371-375.
- Becker AB, Simons FE. Formoterol, a new long-acting selective beta<sub>2</sub>-adrenergic receptor agonist: double-blind comparison with salbutamol and placebo in children with asthma. *J Allergy Clin Immunol* 1989; 84:891-895.
- Bertelsen A, Andersen JB, Busch P, et al. Nebulized sodium cromoglycate in the treatment of wheezy bronchitis: a multicentre double-blind placebo controlled study. *Allergy* 1986; 41:266-270.
- Bisgaard H, Munck SL, Nielsen JP, Petersen W, Ohlsson SV. Inhaled budesonide for treatment of recurrent wheezing in early childhood. *Lancet* 1990; 336:649-651.
- Bisgaard H, Nielsen MD, Anderson B, et al. Adrenal function in children with bronchial asthma treated with beclomethasone dipropionate or budesonide. *J Allergy Clin Immunol* 1988; 81:1088-1095.
- Bisgaard H. Use of inhaled corticosteroids in pediatric asthma. *Ped Pulmonol* 1997; 15:27-33.
- Brogden RN, Sorkin EM. Nedocromil sodium. An updated review of its pharmacological properties and therapeutic efficacy in asthma. *Drugs* 1993; 45:693-715.
- Bronsky EA, Kemp JP, Zhang J, Guerreiro D, Reiss TF. Dose-related protection of exercise bronchoconstriction by montelukast, a cysteinyl leukotriene-receptor antagonist, at the end of a once-daily dosing interval. *Clin Pharm Thera* 1997; 62:556-561.
- Brown PH, Blundell G, Greening AP, Crompton GK. Do large volume spacer devices reduce the systemic effects of high dose inhaled corticosteroids? *Thorax* 1990; 45:736-739.
- Busse WW. The role of leukotrienes in asthma and allergic rhinitis. *Clin Exp Allergy* 1996; 26:868-879.
- Calpin C, Macarthur C, Stephens D, Feldman W, Parkin PC. Effectiveness of prophylactic inhaled steroids in childhood asthma: a systematic review of the literature. *J Allergy Clin Immunol* 1997; 100:452-457.
- Castle W, Fuller R, Hall J, Palmer J. Serevent nationwide surveillance study: comparison of salmeterol with salbutamol in asthmatic patients who require regular bronchodilator treatment. *Brit Med J* 1993; 306:1034-1037.
- Centers for Disease Control and Prevention. General recommendations on immunization. *Morb Mortal Wkly Rep* 1994; Jan 28; 43(RR-1):1-38.
- Chapman KR, Verbeek PR, White JG, Rebuck AS. Effect of a short course of prednisone in the prevention of early relapse after the emergency room treatment of acute asthma. *N Engl J Med* 1991; 324:788-794.
- Check WA, Kaliner MA. Pharmacology and pharmacokinetics of topical corticosteroid derivatives used for asthma therapy. *Am Rev Respir Dis* 1990; 141:S44-S51.

- Clark B. General pharmacology, pharmacokinetics, and toxicology of nedocromil sodium. *J Allergy Clin Immunol* 1993; 92:200-202.
- Clark CE, Ferguson AD, Siddorn JA. Respiratory arrests in young asthmatics on salmeterol. *Respir Med* 1993; 87(3):227-228.
- Clark DJ, Clark RA, Lipworth BJ. Adrenal suppression with inhaled budesonide and fluticasone propionate given by large volume spacer to asthmatic children. *Thorax* 1996; 51:941-943.
- Clissold SP, Heel RC. Budesonide: A preliminary review of its pharmacodynamic properties and therapeutic efficacy in asthma and rhinitis. *Drugs* 1984; 28:485-518.
- Cockcroft DW, Murdock KY. Comparative effects of inhaled salbutamol, sodium cromoglycate, and beclomethasone dipropionate on allergen-induced early asthmatic responses, late asthmatic responses, and increased bronchial responsiveness to histamine. *J Allergy Clin Immunol* 1987; 79:734-740.
- Cockcroft DW, O'Byrne PM, Swystun VA, Bhagat R. Regular use of inhaled albuterol and the allergen induced late asthmatic response. *J Allergy Clin Immunol* 1995; 96:44-49.
- Connett GJ, Warde C, Wooler E, Lenney W. Use of budesonide in severe asthmatics aged 1-3 years. *Arch Dis Child* 1993; 69:351-355.
- Creticos P, Burk J, Smith L, et al. The use of twice daily nedocromil sodium in the treatment of asthma. *J Allergy Clin Immunol* 1995; 95:829-836.
- Crompton G, Duncan J. Clinical assessment of a new breath-actuated inhaler. *Practitioner* 1989; 233:268-269.
- D'Alonzo GE, Nathan RA, Henochowicz S, et al. Salmeterol xinafoate as maintenance therapy compared with albuterol in patients with asthma. *J Am Med Assoc* 1994; 271:1412-1416.
- Dahlen B, Zetterstrom O, Bjorck T, Dahlen SE. The leukotriene-antagonist IC<sub>204</sub>, 219 inhibits the early airway reaction to cumulative bronchial challenge with allergen in atopic asthmatics. *Eur Respir J* 1994; 7:324-331.
- de Benedictis FM, Tuteri G, Pazzelli P, et al. Cromolyn versus nedocromil: duration of action in exercise-induced asthma in children. *J Allergy Clin Immunol* 1995; 96:510-514.
- Djukanovic R, Wilson TW, Britten KM, et al. Effect of an inhaled corticosteroid on airway inflammation and symptoms of asthma. *Am Rev Respir Dis* 1992; 145:669-674.
- Doull IJM, Freezer NJ, Holgate ST. Growth of prepubertal children with mild asthma treated with inhaled beclomethasone dipropionate. *Am J Respir Crit Care Med* 1995; 151:1717-1719.
- Drazen JM, Israel E. Treatment of chronic stable asthma with drugs active on the 5-lipoxygenase pathway. *Int Arch Allergy Immunol* 1995; 107:319-320.
- Eady RP. The pharmacology of nedocromil sodium. *Eur J Respir Dis* 1986; 147:S112-S119.
- Eisen SA, Miller DK, Woodward RS, Spitznagel E, Przybeck TR. The effect of prescribed daily dose frequency on patient medication compliance. *Arch Intern Med* 1990; 150(9):1881-1884.
- Estelle F, Simmons R. A comparison of beclomethasone, salmeterol, and placebo in children with asthma. *N Engl J Med* 1997; 337:1659-1665.
- Everard ML, Clark AR, Milner AD. Drug delivery from holding chambers with attached facemask. *Arch Dis Child* 1992; 67:580-585.
- Fabbri LM, Piattella M, Caramori G, Ciaccia A. Oral vs inhaled asthma therapy: pros, cons and combinations. *Drugs* 1996; 52(S6):20-28.
- Fiocchi A, Riva E, Santini I, et al. Effect of nedocromil sodium on bronchial hyperreactivity in children with nonatopic asthma. *Ann Allergy Asthma Immunol* 1997; 79:503-506.
- Fish JE, Kemp JP, Lockey RF, et al. Zafirlukast for symptomatic mild-to-moderate asthma: a 13-week multicenter study. The Zafirlukast Trialists Group. *Clin Ther* 1997; 19:675-690.
- Fitzpatrick MF, Mackay T, Driver H, Douglas NJ. Salmeterol in nocturnal asthma: a double blind, placebo controlled trial of a long acting inhaled beta<sub>2</sub>-agonist. *Brit Med J* 1990; 301:1365-1368.
- Fuglsand G, Pedersen S. Comparison of Nebuhaler and nebulizer treatment of acute severe asthma in children. *Eur J Respir Dis* 1986; 69:109-113.
- Gaddy JN, Margolskee DJ, Bush RK, Williams VC, Busse WW. Bronchodilation with a potent and selective leukotriene D<sub>4</sub> (LTD<sub>4</sub>) receptor antagonist (MK-571) in patients with asthma. *Am Rev Respir Dis* 1992; 146(2): 358-363.
- Geller M. Acute management of severe childhood asthma. *AACN Clin Issues* 1996; 7:519-0528.
- Geller-Berstein C, Sneh N. The management of bronchial asthma in children under the age of 3 years using Intal (sodium cromoglycate) administered by Spinhaler. *Clin Allergy* 1980; 10:503-508.
- Gemou-Engesaeth V, Bush A, Kay AB, Hamid Q, Corrigan CJ. Inhaled glucocorticoid therapy of childhood asthma is associated with reduced peripheral blood t-cell activation and 'TH<sub>1</sub>-type' cytokine mRNA expression. *Pediatrics* 1997; 99:695-703.
- Glass J, Archer LN, Adams W, Simpson H. Nebulised cromoglycate, theophylline, and placebo in preschool asthmatic children. *Arch Dis Child* 1981; 56:648-651.

Gleeson JG, Price JF. Controlled trial of budesonide given by the nebulizer in preschool children with asthma. *Brit Med J* 1988; 297:163-166.

Goldstein DE, Konig P. Effect of inhaled beclomethasone dipropionate on hypothalamic-pituitary-adrenal axis function in children with asthma. *Pediatrics* 1983; 72:60-64.

Gonzalez JP, Brogden RN. Nedocromil sodium. A preliminary review of its pharmacodynamic and pharmacokinetic properties, and therapeutic efficacy in the treatment of reversible obstructive airways disease. *Drugs* 1987; 34:560-577.

Goodman DC, Littenberg B, O'Connor GT, Brooks JG. Theophylline in acute childhood asthma: a meta-analysis of its efficacy. *Ped Pulmonol* 1996; 21:211-218.

Goodwin A. Asthma therapy-Treating childhood asthma with nebulised steroids. *Professional Care of Mother & Child* 1995; 5:7-9.

Goren A, Noviski N, Avital A, et al. Assessment of the ability of young children to use a powder inhaler device (Turbuhaler). *Pediatr Pulmonol* 1994; 18:77-80.

Green CP, Price JF. Prevention of exercise induced asthma by inhaled salmeterol xinafoate. *Arch Dis Child* 1992; 67: 1014-1017.

Gross NJ. Ipratropium bromide. *N Engl J Med* 1988; 319: 486-494.

Gustafsson P, Tsanakas J, Gold M, et al. Comparison of the efficacy and safety of inhaled fluticasone 200 mcg/day with inhaled beclomethasone dipropionate 400 mcg/day in mild and moderate asthma. *Arch Dis Child* 1993; 69:206-211.

Hahtela T, Jarvinen M, Kava T, et al. Comparison of a beta - agonist, terbutaline, with an inhaled corticosteroid, budesonide, in newly detected asthma. *N Engl J Med* 1991; 325:388-392.

Hahtela T, Jarvinen M, Kava T, et al. Effects of reducing or discontinuing inhaled budesonide in patients with mild asthma. *N Engl J Med* 1994; 331:700-705.

Hanania NA, Wittman R, Kesten S, Chapman KR. Medical personnel's knowledge of and ability to use inhaling devices. Metered-dose inhalers, spacing chambers, and breath-actuated dry powder inhalers. *Chest* 1994; 105(1):111-116.

Hansen-Flaschen J, Schotland H. New treatments for exercise-induced asthma. *N Engl J Med* 1998; 339:129-3.

Harding SM. The human pharmacology of fluticasone propionate. *Respir Med* 1990; 84(Suppl A):25-29.

Hendeles L, Harman E, Huang D, et al. Theophylline attenuation of airway responses to allergen: comparison with cromolyn metered-dose inhaler. *J Allergy Clin Immunol* 1995; 95:505-514.

Henderson WR Jr. The role of leukotrienes in inflammation. *Ann Intern Med* 1994; 121:684-697.

Henriksen JM, Agertoft L, Pedersen S. Protective effect and duration of action of inhaled formoterol and salbutamol on exercise-induced asthma in children. *J Allergy Clin Immunol* 1992; 89:1176-1182.

Higgins RM, Cookson W, Lane DJ, et al. Cardiac arrhythmias caused by nebulized beta-agonist therapy. *Lancet* 1987; 2:863-864.

Hilman BC, Bairnsfather L, Washburne W, Vekovius AL. Nebulized cromolyn sodium: safety, efficacy, and role in the management of childhood asthma. *Ped Allergy Immunol* 1987; 1:43-52.

Holgate ST, Bradding P, Sampson AP. Leukotriene antagonists and synthesis inhibitors: new directions in asthma therapy. *J Allergy Clin Immunol* 1996; 98:1-13.

Holgate ST, Frew AJ. Choosing therapy for childhood asthma. *N Engl J Med* 1997; 337:1690-1692.

Horwitz RJ, McGill KA, Busse WW. The role of leukotriene modifiers in the treatment of asthma. *Am J Respir Crit Care Med* 1998;157:1363-1371.

Ilangovan P, Pedersen S, Godfrey S, Nikander K, Noviski N, Warner JO. Treatment of severe steroid dependent preschool asthma with nebulised budesonide suspension. *Arch Dis Child* 1993; 68:356-359.

Ind PW. Anti-leukotriene intervention: is there adequate information for clinical use in asthma? *Respir Med* 1996; 90:575-586.

Jones AH, Langdon CG, Lee PS, et al. Pulmicort Turbuhaler once daily as initial prophylactic therapy for asthma. *Respir Med* 1994; 88:293-299.

Kamada AK, Hill MR, Ikle DN, Brenner AM, Szeffler SJ. Efficacy and safety of low-dose troleandomycin therapy in children with severe, steroid-requiring asthma. *J Allergy Clin Immunol* 1993; 91:873-882.

Kamada AK, Szeffler SJ. Glucocorticoids and growth in asthmatic children. *Pediatr Allergy Immunol* 1995; 6:145-154.

Kasper WJ, Howe PM. Fatal varicella after a single course of corticosteroids. *Pediatr Infect Dis J* 1990; 9:729-732.

Kelly HW. Asthma pharmacotherapy: current practices and outlook. *Pharmacotherapy* 1997; 17(1 Pt 2):S13-S21.

Kemp JP, Dockhorn RJ, Busse WW, Bleecker ER, Van As A. Prolonged effect of inhaled salmeterol against exercise-induced bronchospasm. *Am J Respir Crit Care Med* 1994; 150: 1612-1615.



- Kemp JP, Furukawa CT, Bronsky EA, et al. Albuterol treatment for children with asthma: a comparison of inhaled powder and aerosol. *J Allergy Clin Immunol* 1989; 83:697-702.
- Kerem E, Levison H, Schuh S, et al. Efficacy of albuterol administered by nebulizer versus spacer device in children with acute asthma. *J Pediatr* 1993; 123:313-317.
- Kerrebijn KF, van Essen-Zandvliet EE, Neijens HJ. Effect of long-term treatment with inhaled corticosteroids and beta-agonists on the bronchial responsiveness in children with asthma. *J Allergy Clin Immunol* 1987; 79(4):653-659.
- Kerstjens HA, Brand PL, Hughes MD, et al. A comparison of bronchodilator therapy for obstructive airways disease. *N Engl J Med* 1992; 327:1413-1419.
- Kim CS, Eldridge MA, Sackner MA. Oropharyngeal deposition and delivery aspects of metered-dose inhaler aerosols. *Am Rev Respir Dis* 1987; 135:157-164.
- Konig P, Eigen H, Ellis MH, et al. The effect of nedocromil sodium on childhood asthma during the viral season. *Am J Respir Crit Care Med* 1995; 152:1879-1886.
- Konig P, Shaffer J. The effect of drug therapy on long-term outcome of childhood asthma: a possible preview of the international guidelines. *J Allergy Clin Immunol* 1996; 98:1103-1111.
- Laitinen LA, Laitinen A, Haahtela T. A comparative study of the effects of an inhaled corticosteroid, budesonide, and a beta-agonist, terbutaline, on airway inflammation in newly diagnosed asthma: a randomized, double-blind, parallel-group controlled trial. *J Allergy Clin Immunol* 1992; 90:32-42.
- Laitinen LA, Laitinen A, Heino M, Haahtela T. Eosinophilic airway inflammation during exacerbation of asthma and its treatment with inhaled corticosteroid. *Am Rev Respir Dis* 1991; 143(2):423-427.
- Lantner RR, Ros SP. Emergency management of asthma in children: impact of NIH guidelines. *Ann Allergy Asthma Immunology* 1995; 74:188-191.
- Leff JA, Busse WW, Pearlman D, et al. Montelukast, a leukotriene-receptor antagonist, for the treatment of mild asthma and exercise-induced bronchoconstriction. *N Engl J Med* 1998; 339:147-52.
- Lemanske RF, Allen DB. Choosing a long-term controller medication in childhood asthma. The proverbial two-edged sword. *Am J Respir Crit Care Med* 1997; 156:685-687.
- Lenney W, Pedersen S, Boner AL, et al. Efficacy and safety of salmeterol in childhood asthma. *Eur J Pediatr* 1995; 154:983-990.
- Leversha AM, Asher MI. Update on spacer devices in childhood asthma. *N Zealand Med J* 1996; 109:76-78.
- Levy J, Zalkinder I, Kuperman O, et al. Effect of prolonged use of inhaler use of inhaled steroids on the cellular immunity of children with asthma. *J Allergy Clin Immunol* 1995; 95:806-812.
- Li JTC, Reed C. Do recent findings warrant a change in approach? Inhaled beta-agonists: weighing risks and benefits. *J Respir Dis* 1993; 14(9):991-1009.
- Lipworth BJ. New perspectives on inhaled drug delivery and systemic bioactivity. *Thorax* 1995; 50:105-110.
- Liu MC, Dube LM, Lancaster J. Acute and chronic effects of a 5-lipoxygenase inhibitor in asthma: a 6-month randomized multicenter trial. *J Allergy Clin Immunol* 1996; 98:859-871.
- Lundback B, Rawlinson DW, Palmer JB. Twelve-month comparison of salmeterol and salbutamol as dry powder formulations in asthmatic patients. European Study Group. *Thorax* 1993; 48(2):148-153.
- McFadden ER Jr, Gilbert IA. Exercise-induced asthma. *N Engl J Med* 1994; 330:1362-1367.
- McGill KA, Busse WW. Zileuton. *Lancet* 1996; 348:519-524.
- McGill KA, Joseph B, Busse WW. Corticosteroids in the treatment of asthma. Practical recommendations. *Clin Immunother* 1995; 4:16-48.
- Meltzer SS, Hasday JD, Cohn J, Bleecker ER. Inhibition of exercise-induced bronchospasm by zileuton: a 5-lipoxygenase inhibitor. *Am J Respir Crit Care Med* 1996; 153:931-935.
- Miraglia del Giudice, Capristo A, Maiello, Apuzzo G. Nebulized sodium cromoglycate for the treatment of asthma in children under five years of age. *Mod Probl Paediatr* 1982; 21:122-127.
- Mollman H, Rohdewald P, Schmidt EW, Salomon V, Derendorf H. Pharmacokinetics of triamcinolone acetonide and its phosphate ester. *Eur J Clin Pharmacol* 1985; 29:85-89.
- Mullen ML, Mullen B, Carey M. The association between beta<sub>2</sub>-agonist use and death from asthma. *J Am Med Assoc* 1993; 270:1842-1845.
- Munoz NM, Douglass I, Mayer D, et al. Eosinophil chemotaxis inhibited by 5-lipoxygenase blockade and leukotriene receptor antagonism. *Am J Respir Crit Care Med* 1997; 155:1398-1403.
- Nassif EG, Weinberger M, Thompson R, Huntley W. The value of maintenance theophylline in steroid dependent asthma. *N Engl J Med* 1981; 304:71-75.
- Nastasi KJ, Heinly TL, Blaiss MS. Exercise-induced asthma and the athlete. *J Asthma* 1995; 32:249-257.
- Nathan RA. Anti-leukotriene agents: a new direction in asthma therapy. *J Asthma* 1996; 33(6):353-366.
- Nelson HS, Hamilos DL, Corsello PR, et al. A double-blind study of troleandomycin and methylprednisolone in asthmatic subjects who require daily corticosteroids. *Am Rev Respir Dis* 1993; 147:398-404.
- Nelson HS, Weber RW. Endocrine aspects of allergic diseases. In: Bierman CW, Pearlman DS, eds. *Allergic diseases from infancy to adulthood*. Philadelphia: WB Saunders, 1988; 15.

- Nelson JA, Strauss L, Skowronski M, et al. Effect of long-term salmeterol treatment on exercise-induced asthma. *N Engl J Med* 1998; 339:141-146.
- Newhouse MT, Dolovich MB. Control of asthma by aerosols. *N Engl J Med* 1986; 315:870-874.
- Newhouse MT, Ruffin RE. Deposition and fate of aerosolized drugs. *Chest* 1978; 73(S6):936-943.
- Newman SP, Millar AB, Lennard-Jones TR, Moren F, Clarke SW. Improvement of pressurized aerosol deposition with Nebuhaler spacer device. *Thorax* 1984; 39:935-941.
- Newman SP, Moren F, Pavla D, Little F, Clarke SW. Deposition of pressurized suspension aerosols inhaled through extension devices. *Am Rev Respir Dis* 1981; 124:317-320.
- Newman SP, Weisz AWB, Talaee N, Clarke SW. Improvement of drug delivery with a breath actuated pressurised aerosol for patients with poor inhaler technique. *Thorax* 1991; 46:712-716.
- Ninan TK, Russell G. Asthma, inhaled corticosteroid treatment, and growth. *Arch Dis Child* 1992; 67(6):703-705.
- Noonan M, Chervinsky P, Busse WW, et al. Fluticasone propionate reduces oral prednisone use while it improves asthma control and quality of life. *Am J Respir Crit Care Med* 1995; 152:1467-1473.
- Novembre G, Frongia GF, Veneruso G, Vierucci A. Inhibition of exercise-induced asthma (EIA) by nedocromil sodium and sodium cromoglycate in children. *Pediatr Allergy Immunol* 1994; 5:107-110.
- O'Byrne PM. Exercise-induced bronchoconstriction: elucidating the roles of leukotrienes and prostaglandins. *Pharmacotherapy* 1997; 17(1 pt 2):31S-38S.
- O'Callaghan C, Cant M, Robertson C. Delivery of beclomethasone dipropionate from a spacer device: what dose is available for inhalation? *Thorax* 1994; 49:961-964.
- O'Hickey SP, Rees PJ. High-dose nedocromil sodium as an addition to inhaled corticosteroids in the treatment of asthma. *Respir Med* 1994; 88:499-502.
- Oliveira CAA, Solé D, Naspitz CK, Rachelefsky GS. Improvement of bronchial hyperresponsiveness in asthmatic children treated for concomitant sinusitis. *Ann Allergy Asthma Immunol* 1997; 79:70-74.
- Pachter LM, Cloutier MM, Bernstein BA. Ethnomedical (folk) remedies for childhood asthma in a main-land Puerto Rican community. *Arch Pediatr Adolesc Med* 1995; 149:982-988.
- Pauwels RA, Joos GF, Kips JC. Leukotrienes as therapeutic target in asthma. *Allergy* 1995; 50:615-622.
- Pauwels RA. New aspects of the therapeutic potential of theophylline in asthma. *J Allergy Clin Immunol* 1989; 83:548-553.
- Pearlman DS, Chervinsky P, LaForce C, et al. A comparison of salmeterol with albuterol in the treatment of mild-to-moderate asthma. *N Engl J Med* 1992; 327:1420-1425.
- Pedersen S, Fuglsang G. Urine cortisol excretion in children treated with high doses of inhaled corticosteroids: a comparison of budesonide and beclomethasone. *Eur Respir J* 1988; 1:433-435.
- Pedersen S, Hansen OR, Fuglsang G. Influence of inspiratory flow rate upon the effect of a Turbuhaler. *Arch Dis Child* 1990; 65:308-310.
- Pedersen S, Hansen OR. Budesonide treatment of moderate and severe asthma in children: a dose-response study. *J Allergy Clin Immunol* 1995; 95:29-33.
- Pedersen S, Mortensen S. Use of different inhalation devices in children. *Lung* 1990; 168:S653-S657.
- Petersen W, Karup-Pedersen F, Friis B, et al. Sodium cromoglycate as a replacement for inhaled corticosteroids in mild-to-moderate childhood asthma. *Allergy* 1996; 51:870-875.
- Pincus DJ, Szeffler SJ, Ackerson LM, Martin RJ. Chronotherapy of asthma with inhaled steroids: the effect of dosage timing on drug efficacy. *J Allergy Clin Immunol* 1995; 95:1172-1178.
- Prahl P, Jensen T, Bjerregaard-Andersen H. Adrenocortical function in children on high-dose steroid aerosol therapy. *Allergy* 1987; 42:541-544.
- Prahl P, Jensen T. Decreased adrenocortical suppression utilizing the nebulizer for inhalation of steroid aerosols. *Clin Allergy* 1987; 17:393-398.
- Prahl P. Adrenocortical suppression following treatment with beclomethasone dipropionate and budesonide. *Clin Exp Allergy* 1991; 21:145-146.
- Ramage L, Lipworth BJ, Ingram CG, Cree IA, Dhillon DP. Reduced protection against exercise induced bronchoconstriction after chronic dosing with salmeterol. *Respir Med* 1994; 88(5):363-368.
- Reisman J, Galdes-Sebalt M, Kazim F, Canny G, Levison H. Frequent administration by inhalation of salbutamol and ipratropium bromide, in the initial management of severe acute asthma in children. *J Allergy Clin Immunol* 1988; 81:16-20.
- Reiss TF, Altman LC, Chervinsky P, et al. Effects of montelukast (MK-0476); a new potent cysteinyl leukotriene (LTD4) receptor antagonist, in patients with chronic asthma. *J Allergy Clin Immunol* 1996; 98:528-534.
- Reiss TF, Sorkness CA, Stricker W, et al. Effects of montelukast (MK-0476) - a potent cysteinyl leukotriene receptor antagonist on bronchodilation in asthmatic subjects treated with and without inhaled corticosteroids. *Thorax* 1997; 52:45-48.
- Risser AL, Mazur LJ. Use of folk remedies in a Hispanic population. *Arch Pediatr Adolesc Med* 1995; 149:978-981.

- Rooklin AR, Lampert SI, Jaeger EA, McGeady SJ, Mansmann HC Jr. Posterior subcapsular cataracts in steroid-requiring asthmatic children. *J Allergy Clin Immunol* 1979; 63(6):383-386.
- Rubin BK, Albers GM. Use of anticholinergic bronchodilation in children. *Am J Med* 1996; 100:49s-53s.
- Ruggins NR, Milner AD, Swarbrick A. An assessment of a new breath-actuated inhaler device in acutely wheezy children. *Arch Dis Child* 1993; 68:477-480.
- Scalabrin DM, Naspitz CK. Efficacy and side effects of salbutamol in acute asthma in children: comparison of oral route and two different nebulizer systems. *J Asthma* 1993; 30:51-59.
- Scarfone RJ, Fuchs SM, Nager AL, Shane SA. Controlled trial of oral prednisone in the emergency department treatment of children with acute asthma. *Pediatrics* 1993; 2:513-518.
- Schnecker MH, Wilson AF, Mukai DS, et al. A device for overcoming discoordination with metered-dose inhalers. *J Allergy Clin Immunol* 1993; 92:783-789.
- Schuh S, Johnson DW, Callahan S, Canny G, Levison H. Efficacy of frequent nebulized ipratropium bromide added to frequent high-dose albuterol therapy in severe childhood asthma. *J Pediatr* 1995; 126:639-645.
- Schwartz HJ, Petty T, Dube LM, et al. A randomized controlled trial comparing zileuton with theophylline in moderate asthma. *Arch Int Med* 1998; 158:141-148.
- Sears MR. The relationship between beta-agonists and asthma mortality. *J Asthma Allergy Pediatr* 1993; 6:123-128.
- Selroos O, Halme M. Effect of a volumatic spacer and mouth rinsing on systemic absorption of inhaled corticosteroids from a metered dose inhaler and dry powder inhaler. *Thorax* 1991; 46:891-894.
- Shapiro GG, Konig P. Cromolyn sodium: a review. *Pharmacotherapy* 1985; 5:156-170.
- Shaw NJ, Edmunds AT. Inhaled beclomethasone and oral candidiasis. *Arch Dis Child* 1986; 61:788-790.
- Silverman M, Connolly NM, Balfour-Lynn L, Godfrey S. Long-term trial of disodium cromoglycate and isoprenaline in children with asthma. *Brit Med J* 1972; 3:378-381.
- Silverstein MD, Yunginger JW, Reed CE, et al. Attained adult height after childhood asthma: effect of glucocorticoid therapy. *J Allergy Clin Immunol* 1997; 99:466-474.
- Simons FE, Persaud MP, Gillespie CA, Cheang M, Shuckett EP. Absence of posterior subcapsular cataracts in young patients treated with inhaled glucocorticoids. *Lancet* 1993; 342: 776-778.
- Sorkness CA. The use of 5-lipoxygenase inhibitors and leukotriene receptor antagonists in the treatment of chronic asthma. *Pharmacotherapy* 1997; 17:50S-54S.
- Spector SL, Smith LJ, Glass M. Effects of 6 weeks of therapy with oral doses of ICI 204, 219, a leukotriene D<sub>4</sub> receptor antagonist, in subjects with bronchial asthma. *Am J Respir Crit Care Med* 1994; 150:618-623.
- Spitzer WO, Suissa S, Ernst P, et al. The use of beta-agonist and the risk of death and near death from asthma. *N Engl J Med* 1992; 326:501-506.
- Stoloff SW. The changing role of theophylline in pediatric asthma. *Am Fam Phys* 1994; 49:839-844.
- Storms WW, Bodman SF, Nathan RA, et al. Use of ipratropium bromide in asthma. *Am J Med* 1986; 81:61-66.
- Strauss RE, Wertheim DL, Bonagura VR, Valacer DJ. Aminophylline therapy does not improve outcome and increases adverse effects in children hospitalized with acute asthmatic exacerbations. *Pediatrics* 1994; 93:205-210.
- Strunk RC, Mrazek DA, Wolfson-Fuhrman GS, LaBrecque JF. Physiological and psychological characteristics associated with deaths due to asthma in childhood. A case-controlled study. *J Am Med Assoc* 1985; 254:1193-1198.
- Suissa S, Dennis R, Ernst P, Sheehy O, Wood-Dauphinee S. Effectiveness of the leukotriene receptor antagonist zafirlukast for mild-to-moderate asthma. A randomized, double-blind, placebo-controlled trial. *Ann Intern Med* 1997; 126:177-183.
- Suissa S, Ernst P, Boivin JP, et al. A cohort analysis of excess mortality in asthma and the use of inhaled beta-agonists. *Am J Respir Crit Care Med* 1994; 149:604-610.
- Sullivan P, Bekir S, Jaffar Z, et al. Anti-inflammatory effects of low-dose oral theophylline in atopic asthma. *Lancet* 1994; 343:1006-1008.
- Svensden UG, Jorgensen H. Inhaled nedocromil sodium as additional treatment to high dose inhaled corticosteroids in the management to bronchial asthma. *Eur Respir J* 1991; 4:992-999.
- Szeffler SJ, Bender BG, Jusko WJ, et al. Evolving role of theophylline for treatment of chronic childhood asthma. *J Pediatr* 1995; 127:176-185.
- Szeffler SJ. Glucocorticoid therapy for asthma: clinical pharmacology. *J Allergy Clin Immunol* 1991; 88:147-165.
- Tabachnik E, Zadik Z. Clinical and laboratory: diurnal cortisol secretion during therapy with inhaled beclomethasone dipropionate in children with asthma. *J Pediatr* 1991; 118:294-297.
- Tasche MJA, van der Wouden JC, Uijen JHJM, et al. Randomised placebo-controlled trial of inhaled sodium cromoglycate in 1-4 year old children with moderate asthma. *Lancet* 1997; 350:1060-1064.

- Thorsson L, Edsbacker S, Conradson TB. Lung deposition of budesonide from Turbuhaler is twice that from a pressurized metered-dose inhaler P-MDI. *Eur Respir J* 1994; 7:1839-1844.
- Tinkelman DG, Reed CE, Nelson HS, Offord KP. Aerosol beclomethasone dipropionate compared with theophylline as primary treatment of chronic, mild to moderately severe, asthma in children. *Pediatrics* 1993; 92:64-77.
- Tomac N, Tuncer A, Saraclar Y, Adalioglu G. Efficacy of salmeterol in the treatment of childhood asthma. *Acta Paediatrica Japonica* 1996; 38:498-494.
- Toogood JH, Jennings B, Greenway RW, Chuang L. Candidiasis and dysphonia complicating beclomethasone treatment of asthma. *J Allergy Clin Immunol* 1980; 65:145-153.
- Tullett WM, Tan KM, Wall RT, Patel KR. Dose-response effect of sodium cromoglycate pressurized aerosol in exercise induced asthma. *Thorax* 1985; 40:41-44.
- Turpeinen M, Sorva R, Juntunen-Backman K. Changes in carbohydrate and lipid metabolism in children with asthma inhaling budesonide. *J Allergy Clin Immunol* 1991; 88:384-389.
- Van Beever HP, Stevens WJ. Pharmacotherapy of childhood asthma. An inflammatory disease. *Drugs* 1992; 44:36-46.
- van Essen-Zandvliet EE, Hughes MD, Waalkens HJ, et al. Effects of 22 months of treatment with inhaled corticosteroids and/or beta<sub>2</sub>-agonists on lung function, airway responsiveness, and symptoms in children with asthma. *Am Rev Respir Dis* 1992; 146:547-554.
- Verberne AAPH, Frost C, Duiverman EJ, et al. Addition of salmeterol versus doubling the dose of beclomethasone in children with asthma. *Am J Respir Crit Care Med* 1998; 158:213-219.
- Verberne AAPH, Frost C, Roorda RJ, et al. One-year treatment with salmeterol compared with beclomethasone in children with asthma. *Am J Respir Crit Care Med* 1997; 156:688-695.
- Vidgren M, Kärkkäinen A, Karjalainen P, Nuutinen J, Paronen P. In vitro and in vivo deposition of drug particles from pressurized aerosol and dry powder inhaler. *Drug Devel Indust Pharm* 1983; 14:2649-2665.
- Waalkens HJ, Van Essen-Zandvliet EE, Hughes MD, et al. Cessation of long-term treatment with inhaled corticosteroid (budesonide) in children with asthma results in deterioration. *Am Rev Respir Dis* 1993; 148:1252-1257.
- Weber RW. Role of long-acting beta<sub>2</sub>-agonists in asthma. *Ann Allergy* 1992; 69:381-384.
- Wechsler ME, Garpestad E, Flier SR, et al. Theophylline in asthma. *N Engl J Med* 1996; 334:1380-1388.
- Wennergren G, Sigurdur K, Strannegård IL. Decrease in hospitalization for treatment of childhood asthma with increased use of antiinflammatory treatment, despite an increase in the prevalence of asthma. *J Allergy Clin Immunol* 1996; 97:742-748.
- Wenzel SE, Kamada AK. Zileuton: the first 5-lipoxygenase inhibitor for the treatment of asthma. *Ann Pharmacother* 1996; 30:858-864.
- Wilson NM, Silverman M. Treatment of acute, episodic asthma in preschool children using intermittent high dose inhaled steroids at home. *Arch Dis Child* 1990; 65:407-410.
- Wolthers OD. Long-, intermediate- and short-term growth studies in asthmatic children treated with inhaled glucocorticosteroids. *Eur Respir J* 1996; 9:821-827.
- Woolcock A, Lundback B, Ringdal N, Jacques LA. Comparison of addition of salmeterol to inhaled steroids with doubling of the dose of inhaled steroid. *Am J Respir Crit Care Med* 1996; 153:1481-1488.
- Woolley M, Anderson SD, Quigley BM. Duration of terbutaline sulfate and cromolyn sodium alone and in combination on exercise-induced asthma. *Chest* 1990; 97:39-45.
- Wright RO, Santucci KA, Jay GD, Steels DW. Evaluation of pre- and post-treatment pulse oximetry in acute childhood asthma. *Acad Emerg Med* 1997; 4:114-117.
- Wurthwein G, Rohdewald P. Activation of beclomethasone dipropionate by hydrolysis to beclomethasone-17-monopropionate. *Biopharm Drug Dispos* 1990; 11:381-394.
- Yates DH, Sussman HS, Shaw MJ, Barnes PJ, Chung KF. Regular formoterol treatment in mild asthma. Effect on bronchial responsiveness during and after treatment. *Am J Respir Crit Care Med* 1995; 152:1170-1174.
- Zachary CT, Evans R III. Perioperative management for childhood asthma. *Ann Allergy Asthma Immunol.* 1996; 77:468-474.
- Zhao JJ, Rogers JD, Holland SD, et al. Pharmacokinetics and bioavailability of montelukast sodium (MK-0476) in healthy young and elderly volunteers. *Biopharm Drug Dispos* 1997; 18:796-777.
- Ziment I, Stein M. Inappropriate and unusual remedies. In: Weiss EB, Stein M, eds. *Bronchial Asthma*. Boston: Little, Brown and Company, 1993; 1145-1151.